



BOULDER MEDICAL CENTER, P.C.

Department of Surgery

Brian Smith, DO
80 Health Park Dr, Suite 100
Louisville, CO 80027
303-666-2632

Christopher Pohlman, M.D.
4745 Arapahoe, Suite 200
Boulder, CO 80303
303-938-4710

Axillary Lymph Node Removal

While recovering from your surgery, please observe the following instructions:

1. Advance your diet slowly as tolerated. Drink plenty of fluids; at least six to eight 8oz glasses of water per day.
2. Take your pain medication as prescribed. You may use ibuprofen, 400-600mg every 8 hours, in addition to your prescribed medication as needed if you do not have a history of ulcers or sensitivity to anti-inflammatory medications. DO NOT drive while on narcotic pain medication.
3. Narcotic pain medications can be very constipating.
 - a. Use a stool softener such as **Pericolace** or **Senokot** one to two times per day. They are available over the counter.
 - b. Once off narcotic pain medications, you may want to switch to **Citrucel** or **Benefiber**, 1-2 tbsp in 8 oz of juice or water one to two times per day.
 - c. If you are off narcotic pain medications and start to have loose bowel movements, you may discontinue the Pericolace, Senokot, Citrucel or Benefiber if you're taking them.
4. If a drain was placed at the time of your surgery, empty the small bulb attached to your **drain** twice daily and record the output. The drain will be ready to remove when the output is less than 30ml per day for 2 days. Call our office when this threshold is met and we will have you come in for drain removal. Some fluid may leak around the drain—this is messy, but not dangerous. This usually means the drain is clogging and needs to be “stripped” to promote drainage.
5. You may remove the dressing over the breast and axillary incisions within 48 hours after surgery and take a shower. Replace the dressing around the drain exit site. **Do not** leave a wet dressing on—it will promote infection. You may leave the dressing on until your first post-op visit and sponge bath if the dressing remains dry.
6. Limit activity on the operative side to no lifting, pushing or pulling. Do not raise the arm more than 90 degrees. Exercises can generally begin within 48 hours after the drain is removed. We will instruct you in these, or refer you to Physical Therapy when the drain is removed.
7. Please call if you experience fevers, chills, painful enlarging swelling around incisions, redness or streaking around incisions, or persistent bleeding.
8. If you have not already scheduled a post-operative appointment, please call our office to schedule this appointment 7-10 days after surgery. Your final pathology report will usually be available for review at that time.

WS.Surg.18 07/22

2750 Broadway
Boulder, CO 80304

4745 Arapahoe,
Suite 200
Boulder, CO 80303

80 Health Park Dr,
Suite 100
Louisville, CO 80027

1551 Professional Lane
Suite 270
Longmont, CO 80501

80 S. Briggs
Suite 100
Erie, CO 80516