GASTROENTEROLOGY
(303) 440-3223

SUPREP Colonoscopy Procedure Instructions

Please carefully read through the instructions below to prepare for your procedure.
If you have questions, please call us at (303) 440-3223, Mon - Fri 8am - 4:30pm. **DO NOT REPLY TO THIS MESSAGE.**

You will receive an automated text or email communication one week before your procedure prompting you to review the instructions below.

Your procedure will be performed at the Boulder Medical Center Ambulatory Surgery Center located on the 2nd floor at 2750 Broadway in Boulder.

Anesthesia Service
All of our anesthesia services are provided by Physician Anesthesia Services (PAPC). **Please refer to "Anesthesia & Sedation Policy" below in these instructions.** If Boulder Medical Center and your provider are in-network with your insurance company, the anesthesia service should be too. If you have questions about coverage, call (303) 309-0348.

INSTRUCTIONS
- **DO NOT have breakfast the day before the procedure**
- We have sent the laxative prescription to your pharmacy on file. Please follow the instructions below rather than on the prescription package
    - It is very important to follow these instructions to ensure full visualization of your colon
    - Inadequate preparation limits the value of this procedure and could necessitate rescheduling your exam

TIMELINE
- **5 pm the night before your procedure,** take the first dose of prep medication
- **4 hours before procedure**, take the second dose of prep medication
- **3 hours before the procedure, nothing by mouth,** not even water
- Arrival time: 45 minutes before the procedure. Patients with an 8:00 AM procedure should arrive 45 minutes prior to your procedure.
- Departure time: 45 minutes after procedure

Day BEFORE Procedure
- **NO SOLID FOODS**
- **For breakfast** you may have one Ensure (or similar- Fairlife, Boost) supplement in chocolate or vanilla flavor the morning prior to the procedure. You may also add a protein powder of your choosing

Boulder Medical Center

Broadway	Avista	Foothills	Longmont	Erie
2750 Broadway Boulder, CO 80304 303-440-3000	*80 Health Park Dr* *Ste 100* *Louisville, CO 80027* *303-673-0448*	*4745 Arapahoe Rd* *Ste 200* *Boulder, CO 80303* *303-938-4700*	*1551 Professional Ln.* *Ste 240* *Longmont, CO 80501* *303-938-4710*	*80 S Briggs Rd Ste 105* *Erie, CO 80516* *Peds 303-673-0448* *FP 303-440-3102*

- Drink **ONLY CLEAR LIQUIDS** the rest of the day
- **5 PM**: Start drinking the first dose of the SUPREP solution.
    - Drink at least 32 ounces of clear liquid afterward. Continue drinking clear liquids all night.
    - If you do NOT start having bowel movements within 2 hours of completing this (5pm) dose, get 2 bottles of Magnesium Citrate or 4-6 doses of Miralax and take them immediately.

Day of Procedure

- **NO SOLID FOODS**
- **4 hours before your procedure**, drink the second portion of the laxative solution. You must complete the second half even if your stools look watery.
- Drink at least 32 ounces of clear liquids afterward.
- If you do NOT start having bowel movements within 2 hours of completing this (morning) dose, get 2 bottles of Magnesium Citrate or 4-6 doses of Miralax and take them immediately.
- After completing the laxative preparation, you MAY continue to drink clear liquids up to 3 hours before your procedure.
- Nothing by mouth 3 hours before your appointment, not even a sip of water.
- Do not consume any food. If you should begin to vomit, stop drinking the preparation for 20-30 minutes and then try finish the laxative preparation. If you are unable to finish the bowel preparation due to vomiting, call (303) 440-3223 and ask to speak to a nurse.
- No smoking.
- No gum, hard candy, or breath mints.
- No jewelry.
- Ensure you have a driver to take you home after the procedure.
- Take your regular medications the day before the procedure, unless otherwise directed by our staff or your doctor. See additional instructions below for more details.
- If you do NOT start having bowel movements within 2 hours of completing that dose, go to a pharmacy and get 2 bottles of Magnesium Citrate or 4-6 doses of Miralax and take them immediately. Repeat this process if you also vomit after the morning dose of the colon prep.

Clear Liquid Diet Items that ARE Allowed

- Water, coconut water (no pulp)
- Broth: (without pulp)
- Apple juice or cider
- Lemonade
- Italian ices, popsicles (no fruit, cream, red or purple)
- Coffee (without cream or milk)
- Clear gelatin (without fruit and not red or purple)
- Tea

Liquids that are NOT Allowed

- Red or purple juices or other fluids
- Milk
- Cream
- Oatmeal
- Cream of Wheat
- Tomato juice
- Orange juice
- Cream soups
- Any soup other than the listed broth
- Grapefruit juice

- Milkshakes
- Alcohol

GENERAL PROCEDURE INSTRUCTIONS
Upon arrival, please check in at the reception desk located in the Gastroenterology waiting room on the second floor. Be sure to bring your insurance card.

Transportation
You will be receiving intravenous (I.V.) anesthesia for your procedure so you may not drive or return to work. An adult must pick you up in person and sign discharge paperwork before you can leave the facility. Rideshare services (Uber, Lyft), taxis, or buses are **not permitted** unless you are accompanied by a friend or family member. Medical transportation services such as Dignity Care are acceptable alternatives.

If you are unable to provide appropriate transportation details at check-in, your procedure will be canceled. To reschedule or cancel your appointment, please call (303) 440-3216.

Recovery
After the procedure, you will recover in the Ambulatory Surgery Center (ASC) recovery unit until the sedation wears off. Most patients are discharged approximately 45 minutes after arrival into the PACU.

In the event a polyp is removed, you may be advised to remain in town for up to three days following the procedure.

Cancel or Reschedule
Cancellations must be made at least four (4) business days before the procedure to avoid a $250 cancellation fee.

If you have medical or preparation instruction questions, contact us at (303) 440-3223.

Medications
Please follow the timeline below for stopping certain medications before your procedure. If you have any questions about stopping a medication, contact your prescribing provider.

Parkinson's medications: Continue usual dosing, **timed with prep** for proper absorption.

When to Stop	Which Medications	Examples
7 Days Before Procedure	GLP-1 / Semaglutide medications	Ozempic, Wegovy, Zepbound, Rybelsus
	Diet pills / stimulants	Phentermine
	Iron supplements	
	Fish oil / omega-3 supplements	
5 Days Before Procedure, if approved by prescribing provider	Blood thinners	Warfarin (Coumadin), Clopidogrel (Plavix), Aspirin
	NSAIDs	Advil, Aleve, Ibuprofen
3 Days Before Procedure	SGLT2 inhibitors (gliflozins)	Jardiance, Farxiga
2 Days Before Procedure	Eliquis	Apixaban
24 Hours Before Procedure	**Half the usual dose** of long-acting insulin	Lantus
	Stimulants	Adderall
	Marijuana in any form	
	Smoking of any kind	
Day of Procedure	Blood pressure medications ending in "-pril" or "-sartan"	Lisinopril, Losartan
	Diuretics ("water pills")	Lasix, Hydrochlorothiazide (HCTZ), Bumex
	Diabetes pills used to lower blood sugar**	Metformin
	Short-acting insulin	Regular / R insulin
	Benzodiazepines	Continue usual dose if taken regularly, but the patient must be able to **consent for anesthesia**

**Contact your prescribing physician before stopping these medications if they were prescribed for a medical condition.

Medications You CAN Take the Morning of Your Procedure

- Thyroid medications
- Anti-epileptic medications
- Long-term steroid medications

Boulder Medical Center

Broadway	Avista	Foothills	Longmont	Erie
2750 Broadway Boulder, CO 80304 303-440-3000	*80 Health Park Dr* *Ste 100* *Louisville, CO 80027* *303-673-0448*	*4745 Arapahoe Rd* *Ste 200* *Boulder, CO 80303* *303-938-4700*	*1551 Professional Ln.* *Ste 240* *Longmont, CO 80501* *303-938-4710*	*80 S Briggs Rd Ste 105* *Erie, CO 80516* *Peds 303-673-0448* *FP 303-440-3102*

- Inhalers (maintenance and rescue)
- PPI's (omeprazole (prilosec), pantoprazole (protonix), esomeprazole (nexium)
- Beta blockers (medications ending in "-olol" such as Metoprolol, Atenolol, Labetalol)

Please refrain from taking all other medications on the day of the procedure unless otherwise instructed by your provider.

About the Procedure

Screening vs Diagnostic Colonoscopy

The colonoscopy itself is mostly the same. However, the REASON for the colonoscopy is different. A screening colonoscopy is like getting mammograms or prostate checks; it is done in everyone at a certain age because your risk for colon cancer increases as you age. Since most colon cancers grow from colon polyps, and colon polyps do not cause symptoms (meaning you would not know you have any polyps), we look in everyone's colon for polyps. If we find polyps, we remove them to prevent those polyps from growing into cancer.

A diagnostic colonoscopy is any colonoscopy that does not qualify for screening. This includes a colonoscopy for any symptom (rectal bleeding, diarrhea, weight loss, Crohn's disease, ulcerative colitis, IBS, etc.) or for a positive colon cancer screening test (Cologuard, FIT, FOBT, CT colonography). The reason for the colonoscopy can change what your doctor does (take biopsies, band hemorrhoids). You will have time to speak with your doctor in the pre-operative area prior to your procedure to go over this.

Insurance Coverage for Screening vs Diagnostic

A screening colonoscopy (CPT code 45378) uses the ICD-10 code Z12.11 and typically will fall under the preventative health portion of your insurance company. Most insurance companies fully cover the cost of preventative health services. There may be a cost to you for the anesthesia services or lab services if your doctor removes polyps/takes biopsies because the procedure then is considered diagnostic.

To qualify for a screening colonoscopy, you must be between the ages of 45-75, have no symptoms or family members with colon cancer, and your last colonoscopy must have been at least 10 years ago. You can call your insurance carrier with the codes listed above to find out how your insurance plan covers this.

A diagnostic colonoscopy (CPT code 45378) can use many different codes. Common ICD-10 codes are: Z80.0 Family history of colon cancer, Z86.010 Personal history of colon polyps, R19.5 Positive Cologuard/FOBT/FIT testing.

If you have had a prior SCREENING colonoscopy and had precancerous polyps removed, any subsequent colonoscopy is typically considered a DIAGNOSTIC colonoscopy by your insurance carrier. If your colonoscopy is being done for any symptom, then it will be considered diagnostic. If your

colonoscopy is being done before the age of 45, then it will be considered diagnostic. If your colonoscopy is done prior to every 10 years, it will most likely be considered diagnostic.

Dealing with Nausea
Nausea is quite common with all the colon preps. To help reduce nausea with a medication you should avoid taking your prep all it at once. We recommend that you complete your prep slowly to prevent nausea as much as possible

No Bowel Movements After Prep
If you do NOT start having bowel movements within 2 hours of completing your 5 pm dose, get 2 bottles of Magnesium Citrate or 4-6 doses of Miralax and take them immediately. If you do NOT start having bowel movements within 2 hours of completing your morning dose, get 2 bottles of Magnesium Citrate or 4-6 doses of Miralax and take them immediately.

Vomiting Prep
As nausea is common with all colon preps, vomiting can be too. Since it is impossible to know how much of the prep you threw up, we recommend you wait for 2 hours after completing the dose of prep to see if you start having bowel movements. If you have several bowel movements within 2 hours of taking the prep, then you have most likely retained enough of the medication to work effectively**.**

If you do NOT start having bowel movements within 2 hours of completing the 5pm dose, get 2 bottles of Magnesium Citrate or 4-6 doses of Miralax and take them immediately. Repeat this process in the morning if you also vomit after the morning dose of the colon prep.

Prep Instructions
Please follow the instructions sent to you from BMC. We **DO NOT** want you to eat breakfast the day before as this tends to cause you to not clean out all the way.

Constipation
If you are typically constipated, the prep may not be as effective. We recommend you prepare early by taking 1 capful of Miralax every day for 3 days prior to your day of clear liquids to help get your bowels moving. A low fiber diet beforehand may be helpful as well.

Costs and Insurance Coverage
Lab Service
If you would like to check your insurance coverage for lab work, our providers use LabCorp pathology lab.

Anesthesia & Sedation Policy
EGD Procedures:
Sedation is required for all EGD procedures. Patients may not decline anesthesia for these cases.

Colonoscopy Procedures:
Patients have the right to decline sedation for a colonoscopy. However, due to scheduling and staffing requirements, all colonoscopies performed without sedation must be scheduled as the **first or last case of the day.**

If you choose to proceed without anesthesia, **you must notify our office at least 48 hours prior to your scheduled procedure** so we can make the appropriate scheduling adjustments. Please call us at 303-440-4217.

Late Changes & Day-of-Procedure Policy:
- If you arrive on the day of your procedure and decline anesthesia without prior notice, your procedure will be rescheduled and a **$250 cancellation fee** will apply.

- If you previously declined anesthesia but decide on the day of your procedure that you would like sedation, your procedure will also be rescheduled and a **$250 cancellation fee will apply** due to the last-minute need to reschedule.

A Note About Cigna and Aetna: Our anesthesia provider, Physician Anesthesia Services, will transition to out-of-network with Cigna Healthcare effective 7/31/2026 and Aetna on 11/15/2026. Under the No Surprises Act, if you are in-network at BMC, Cigna and Aetna should process your anesthesia claim at the in-network benefit level. Please contact Cigna or Aetna if you have questions.

Out of Pocket Cost (without insurance)
- **Facility fee:** $950
- **Physician fee:** ranges from $300-$800 depending on what the physician does during the procedure.
- **Anesthesia fee:** Call Ventra Health for estimates at (303)309-0348
- **Lab services:** The cost to the patient **per specimen** for a GI procedure if not billed through insurance is around $200.

Please contact your insurance carrier if you have questions about in-network labs and charges.

Pre-Certification vs. Covered Benefit vs. Covered Payment
Pre-certification is a requirement for most insurers, notifying them that a medical procedure is being scheduled. This gives them the opportunity to deny the procedure if they choose.

- Pre-certification of this procedure with your insurer will be performed by our office.
- Pre-certification does NOT determine that a medical procedure is a "covered benefit" under your policy. Although your insurance provider may issue a "pre-approval" based on the information submitted during our pre-certification process, this **does not guarantee how much of the procedure will be covered or how much you may be responsible for paying. Final coverage and out-of-pocket costs for GI procedures are often not determined until after the procedure is completed.**

Boulder Medical Center

Broadway	Avista	Foothills	Longmont	Erie
2750 Broadway Boulder, CO 80304 303-440-3000	*80 Health Park Dr* *Ste 100* *Louisville, CO 80027* *303-673-0448*	*4745 Arapahoe Rd* *Ste 200* *Boulder, CO 80303* *303-938-4700*	*1551 Professional Ln.* *Ste 240* *Longmont, CO 80501* *303-938-4710*	*80 S Briggs Rd Ste 105* *Erie, CO 80516* *Peds 303-673-0448* *FP 303-440-3102*

- A procedure initially scheduled as a routine screening may be classified as diagnostic—or vice versa—depending on the findings and actions taken by the physician during the procedure. As a result, we are unable to confirm in advance whether your insurance will categorize your procedure as screening or diagnostic, or what the exact cost to you will be.
- Your procedure will be performed using the services of an anesthesiologist who will bill your insurance for you. You can contact the anesthesia group to verify they are contracted with your insurance. You will receive a separate bill from the anesthesia group after the procedure.
- If you wish to proceed with a test that is not a covered benefit, then you will be asked to sign a waiver accepting personal responsibility for payment.

Medicare

We have Medicare benefit information available, but you may still need to call your secondary carrier. Medicare has very specific rules as to timing of colonoscopy procedures. Please be sure yours will be a "covered benefit" by calling 1-800-MEDICARE.

Contact your insurance carrier if you have questions about coverage and payment.

Boulder Medical Center

Broadway	Avista	Foothills	Longmont	Erie
2750 Broadway Boulder, CO 80304 303-440-3000	*80 Health Park Dr* *Ste 100* *Louisville, CO 80027* *303-673-0448*	*4745 Arapahoe Rd* *Ste 200* *Boulder, CO 80303* *303-938-4700*	*1551 Professional Ln.* *Ste 240* *Longmont, CO 80501* *303-938-4710*	*80 S Briggs Rd Ste 105* *Erie, CO 80516* *Peds 303-673-0448* *FP 303-440-3102*