Department of Surgery

Stephen Johs, M.D. 80 Health Park Dr, Suite 100 Louisville, CO 80027 303-666-2632 Christopher Pohlman, M.D. 2750 Broadway Boulder, CO 80304 303-440-3093 Sandra Taylor, M.D. 4745 Arapahoe, Suite 200 Boulder, CO 80303 303-938-4710

Call 303-440-3000 to reach the on-call surgeon after normal business hours.

Port Placement

While recovering from your surgery, please observe the following instructions:

- 1. Advance your diet slowly as tolerated. Drink plenty of fluids; at least six to eight 8oz glasses of water per day.
- 2. Take your pain medication as prescribed. You may use ibuprofen, 400-600mg every 8 hours, if you do not have a history of ulcers or sensitivity to anti-inflammatory medications.
- 3. You may remove the dressing over the incision within 48 hours after surgery and take a shower. **Do not** leave a wet dressing on—it will promote infection.
- 4. Limit activity on the side your port is on to no pushing or pulling, and avoid contact sports or strenuous exercise while the incision is healing.
- 5. Please call if you experience fevers, chills, painful enlarging swelling around the incision, redness or streaking around the incision, or persistent bleeding. Call your Surgeon's office, or 303-440-3000 for the on-call surgeon after-hours.
- 6. Your port will need to be flushed at least once a month, and after each use. This can be done at your oncologist's office.

Surg.23 04/16